

THE EFFECTS OF RUMINAL ESCAPE PROTEIN OR FAT ON NUTRITIONAL STATUS OF PREGNANT WINTER-GRAZING BEEF COWS¹

J. L. Miner^{2,3}, M. K. Petersen², K. M. Havstad², M. J. McInerney²
and R. A. Bellows⁴

Montana State University, Bozeman 59717

ABSTRACT

This study was designed to determine the influence of soybean meal supplementation, with or without additional ruminal escape protein or fat, on the nutritional status of pregnant winter-grazing beef cows. During two winters (Trials 1 and 2), approximately 60 prepartum beef cows grazed native foothills range each year. Cows were allotted randomly to five groups and supplemented (g/d) with either none (control); 570 soybean meal (SOY); 450 soybean meal plus 230 blood meal (SOY + BM); 140 soybean meal, 16 urea plus 450 corn gluten meal (SOY + CGM); or 570 soybean meal plus 210 animal fat (SOY + FAT). These supplements were designed to supply similar quantities of ruminal degraded protein while varying in escape protein quantity and source (SOY + BM and SOY + CGM). Condition scores and body weights were determined at trial initiation (mid-December) and conclusion (early March). Eight blood samples obtained over 4 d during three periods (9, 4 and 1 wk prior to parturition) were analyzed for concentrations of glucose, urea nitrogen (N), total bilirubin, creatinine, albumin, total protein and cholesterol. Cows in the control treatment experienced the greatest BW loss in both trials. In Trial 2, escape protein tended to decrease ($P < .06$) BW loss compared to SOY, though loss tended to be greater ($P < .08$) with SOY + CGM than with SOY + BM. Escape protein can enhance nutritional status when supplemented with .6 kg/d of soybean meal.

(Key Words: Beef Cattle, Protein Supplements, Blood Chemistry.)

J. Anim. Sci. 1990. 68:1743-1750

Introduction

Pregnant beef cows grazing winter range forage may lose up to .5 kg BW/d (Pinney et al., 1972). Supplementation of grazed forage with .5 to 1 kg of soybean meal or cottonseed meal/d typically reduces prepartum weight loss (Lusby and Wagner, 1987). Supplementation

with nonprotein nitrogen plus cereal grain also can reduce weight loss but its benefit usually is less than that resulting from supplementation with natural protein (Mies et al., 1967; Thomas et al., 1968).

Some of the response to soybean meal or cottonseed meal supplements may be due to ruminal escape protein. The gestating cow, compared to the nonpregnant cow, may have a higher requirement for glucogenic substrates (MacRae and Loble, 1986). Escape protein may fulfill this need via gluconeogenesis. Petersen and Clanton (1981) supplemented winter-grazing pregnant cows with soybean meal or blood meal plus urea. They found that blood meal plus urea was inferior to an isonitrogenous soybean meal supplement as measured by cow BW loss.

This study was designed to determine the influence of soybean meal supplementation,

¹Contribution No. J-2093 from the Montana Agric. Exp. Sta. The authors express their appreciation to Steven Beverlin, Roger Dunn and Eldon Ayers for their excellent technical assistance.

²Anim. and Range Sci. Dept. Address reprint requests to Mark K. Petersen.

³Present Address: Dept. of Anim. Sci., Univ. of Missouri, Columbia 65211.

⁴Fort Keogh Livest. and Range Res. Lab. USDA-ARS, Miles City, MT 59301.

Received January 15, 1989.

Accepted October 19, 1989.

with or without ruminal escape protein or fat, on nutritional status of pregnant, winter-grazing beef cows. Nutritional status was defined as change in BW, condition score and specific blood metabolite concentrations.

Materials and Methods

Animals. Two trials were conducted at the Red Bluff Research Ranch, Norris, MT. Trial 1 began December 15, 1984 and ended March 1, 1985. Trial 2 began December 15, 1985 and ended March 6, 1986. For each trial, pregnant 3- to 6-yr-old crossbred (Angus \times Hereford and Tartentaise) cows (Trial 1, $n = 55$; Trial 2, $n = 61$) were selected randomly within each age from the ranch herd of 150 cows. Those selected were expected to calve within a 21-d period.

Pasture. The pasture used in these two trials was described by Miner and Petersen (1989).

Treatments. Prior to each trial, cows were allotted randomly within each age to five supplement (S) treatments designated as control (no supplement), SOY (570 g soybean meal/d), SOY + BM (450 g soybean meal plus 230 g blood meal/d), SOY + CGM (140 g soybean meal, 16 g urea plus 450 g corn gluten meal/d) and SOY + FAT (570 g soybean meal plus 210 g animal fat/d) (Miner and Petersen, 1989). The control received range forage only. The SOY group and all other protein-supplemented groups received a calculated equivalent of .2 kg/d of ruminally degraded protein. Cows were fed their supplement individually at 48-h intervals at 1300. Other details of the supplement regimen were described by Miner and Petersen (1989).

Measurements

Cow Weights, Condition Scores and Calf Birth Weights. Condition was scored by two technicians (using both palpation and observation) and BW was recorded two consecutive days after 12 h of feed and water deprivation at the start and finish of each trial. Calf sex, birth dates and birth weights also were recorded.

Blood Metabolites. During three periods, 9,

4 and 1 wk prior to the first predicted parturition date (designated as 9 wk, 4 wk and 1 wk, respectively) in each trial, blood samples were obtained from either the jugular vein (Trial 1) or from an artery or vein near the base of the tail (Trial 2). Although most animals were sampled easily from the jugular vein in Trial 1, effects of stress were minimized with the tail bleeding technique because animals were not restrained in a squeeze chute. In Trial 1, nine blood samples were taken from each of four cows per treatment during each period. In Trial 1, samples were obtained -5, 0, 5, 10, 20, 30, 43, 67 and 91 h post-supplementation. In Trial 2, all cows (61) were sampled 0 and 48 h post-supplementation. In addition, four cows per treatment were bled at 5, 10, 20, 30 and 75 h after supplementation. Samples were taken within the 48-h supplementation cycle to account for diurnal variation in blood metabolite concentrations. Samples were limited to four cows per treatment in Trials 1 and 2 during the 48-h supplementation cycle to coincide with another sampling protocol as described by Miner and Petersen (1989). In both trials, two 10-ml samples were obtained, one in an untreated plastic syringe (Trial 1) or vacuum tube⁵ (Trial 2) and one in a syringe (Trial 1) or vacuum tube (Trial 2) containing 2 mg/ml of sodium fluoride and an anticoagulant. Samples in containers with anticoagulant were centrifuged immediately at 500 \times g for 30 min. Plasma was decanted into two 12-mm \times 75-mm polystyrene (Trial 1) or polypropylene (Trial 2) culture tubes and frozen immediately. Other samples were allowed to coagulate for 4 h and centrifuged; serum was decanted and frozen. Plasma was analyzed for glucose concentration; serum was analyzed for concentration of urea N, total bilirubin, creatinine, albumin, cholesterol and total protein⁶. No serum was obtained during period 1 of Trial 1.

Statistical Analyses. Cow BW condition score and calf birth weight were analyzed by ANOVA using the GLM procedure of SAS (1984). Main effects included supplement and cow age. Initial cow BW and condition score were included as linear covariates in the least squares model. Age classes were 3-, 4- and 5- to 6-yr-old cows. Supplement least squares means were separated with single degree of freedom nonorthogonal contrasts. Standard errors were pooled by averaging the standard errors of subclass means. Blood

⁵Becton-Dickerson, Rutherford, NJ.

⁶Technicon Instruments Corp., Tarrytown, NY. Analyses conducted by Marsh Laboratory, Montana State University, Bozeman.

metabolites were analyzed by multivariate analysis of variance utilizing the General Linear Models procedure of SAS (1984). The repeated measures procedure was utilized in order to account for the repeated sampling of the same cow in each 48-h post-supplementation period. Sampling period and supplement were included in the model as main effects. Tests of significance and single degree of freedom linear contrasts employed cow (period) mean square as the error term.

Results

Climate

The average median daily temperature was lower during period 2 of Trial 1 (-6.5°C) than during either period 1 (-4.2°C) or period 3 (1.1°C ; National Oceanic and Atmospheric Administration, 1985). The ambient temperature also was declining with each day during 4 wk of period 2, whereas climatic conditions remained relatively constant during the other two periods. Snow cover in the pasture was greatest during 4 wk.

In Trial 2, the average median daily temperature was warmer than in Trial 1 during sampling periods (9 wk = -2.3°C ; 4 wk = 2.1°C ; 1 wk = 9.2°C ; National Oceanic and Atmospheric Administration, 1986). Snow cover was greater in Trial 2 except during the sampling periods.

Cow Weights, Condition Scores and Calf Birth Weights

Trial 1. Supplement affected ($P < .01$) cow BW change (Table 1). Control cows lost 1.9 kg during the 76-d study, whereas supplemented cows gained an average of 36.3 kg. No differences were detected among the groups fed supplement. Neither cow age group, initial cow weight or initial condition score affected ($P > .1$) BW change.

Supplement affected ($P = .02$) cow condition score change (Table 1). Control cows lost 1.46 units, which was greater ($P < .01$) than the mean loss of the supplemented cows. The condition score loss for the mean of cows fed SOY + BM and SOY + CGM was less ($P = .03$) than that for cows fed SOY alone. Cow age did not ($P > .05$) influence response to supplement. The regression coefficients ($P < .05$) of condition score change on initial BW

TABLE 1. LEAST SQUARES MEANS FOR CALF BIRTH WEIGHT AND PRECALVING COW BODY WEIGHT AND CONDITION SCORE CHANGE (TRIALS 1 AND 2)

Trial and supplement ^a	BW change ^b , kg	Condition score change ^c	Calf birth wt, kg
Trial 1			
Control	-1.9	-1.46	38.4
SOY	31.7	-1.18	38.1
SOY+BM	38.2	-.76	39.2
SOY+CGM	44.4	-.76	37.7
SOY+FAT	30.7	-.56	39.1
SE	7.04	.212	1.03
Trail 2			
Control	-46.4	-.95	37.8
SOY	-20.1	-.93	38.7
SOY+BM	-1.8	-.46	38.4
SOY+CGM	-15.0	-.69	39.2
SOY+FAT	-10.1	-.63	38.0
SE	5.37	.122	1.14

^aSOY = soybean meal, SOY+BM = soybean meal plus blood meal, SOY+CGM = soybean meal plus corn gluten meal, SOY+FAT = soybean meal plus animal fat.

^bTrial 1 and Trial 2: Control vs others, $P < .01$ and $P < .01$, respectively. Trial 2: SOY vs SOY+BM, SOY+CGM, $P = .06$; SOY+BM vs SOY+CGM, $P = .08$.

^cTrial 1 and Trial 2: Control vs others, $P < .01$ and $P = .04$; SOY vs SOY+BM, SOY+CGM, $P = .03$ and $P < .01$ respectively.

and condition score were .003 and $-.35$, respectively.

Neither treatment, cow age group nor initial condition score affected calf birth weight (Table 1). The regression coefficient ($P < .01$) of birth weight on initial cow body weight was .05.

Trial 2. Supplement affected ($P < .01$) cow BW change (Table 1). Control cows lost 46.4 kg during the 81-d study; this was greater ($P < .01$) than the mean weight loss for supplemented cows. Cows fed SOY lost 20.1 kg; this was greater ($P = .06$) than for cows fed SOY + BM (-1.8 kg) and SOY + CGM (-15.0 kg). Cows fed SOY + BM tended to lose less ($P = .08$) weight than cows in SOY + CGM. Neither cow age group, initial cow weight or initial condition score influenced ($P > .05$) weight change.

Supplement affected ($P = .03$) cow condition score change (Table 1). The control lost .95 units, which was more ($P = .04$) than for S groups. The loss for the mean of cows fed SOY + BM and SOY + CGM was less ($P < .01$) than for cows fed SOY. Cow age did not ($P > .05$) influence response to supplement. The regression coefficients ($P < .01$) of

TABLE 2. LEAST SQUARES MEANS FOR PRECALVING CONCENTRATIONS OF GLUCOSE, ALBUMIN, TOTAL PROTEIN, UREA NITROGEN, CREATININE, TOTAL BILIRUBIN AND CHOLESTEROL AS INFLUENCED BY SUPPLEMENT AND WEEK PRIOR TO PARTURITION (TRIAL 1)

Metabolite	Supplement ^a					Week			
	Control	SOY	SOY+ BM	SOY+ CGM	SOY+ FAT	9	4	1	SE
Glucose mg/dl ^{bc}	53.5	58.6	58.0	53.6	59.7	63.7	47.5	58.9	1.3
Albumin g/dl	2.4	2.7	2.9	2.7	2.8		2.8	2.7	.3
Total protein g/dl	7.9	7.8	7.8	7.9	7.5		7.8	7.7	.2
Urea nitrogen mg/dl ^d	7.7	6.3	9.0	9.8	6.2		6.4	9.2	1.2
Creatinine mg/dl ^e	1.9	1.8	1.6	1.6	1.5		1.6	1.8	.1
Total bilirubin ^{f,g}	.13	.12	.06	.09	.10		.12	.08	.03
Cholesterol mg/dl ^h	96.7	113.6	95.3	90.3	142.3		108.8	106.5	7.2

^aSOY = soybean meal, SOY+BM = soybean meal plus blood meal, SOY+CGM = soybean meal plus corn gluten meal, SOY+FAT = Soybean meal plus animal fat.

^bControl vs other supplements, $P < .1$.

^cEffect of week prior to parturition, $P < .01$.

^dSOY vs SOY+BM, SOY+CGM, $P < .05$; SOY+FAT vs SOY+BM, SOY+CGM, $P < .05$.

^eControl vs other supplements, $P < .05$.

^fSOY vs SOY+BM, SOY+CGM, $P < .05$.

^gEffect of week prior to parturition, $P < .05$.

^hSOY+FAT vs SOY+BM, SOY+CGM, $P < .01$.

condition score change on initial weight and condition score were .002 and $-.73$, respectively.

Supplementation did not affect ($P > .05$) calf birth weight (Table 1) in Trial 1 or 2.

Blood Metabolites

Trial 1. Plasma glucose concentration was highest in SOY + FAT and lowest in SOY + CGM and control, with other S intermediate ($P = .06$). All S groups had lower ($P < .01$) plasma glucose concentration at 4 wk than at 9 wk prepartum (Table 2).

Albumin and total protein were unaffected ($P > .05$) by supplement or week prior to parturition. Serum urea nitrogen was higher ($P < .05$) for SBM + BM and SBM + CGM than for SOY or SOY + FAT and higher at 1 wk than at 4 wk prepartum (Table 2).

Creatinine was lower ($P < .05$) in those cattle receiving supplement, whereas total bilirubin was lower in cows receiving SOY + BM or SOY + CGM than in cows receiving SOY (Table 2). Total bilirubin concentration was higher ($P < .05$) and creatinine was lower at 4 wk than at 1 wk prepartum for all S groups. Blood serum cholesterol concentration with SOY + BM and SOY + CGM supplements was lower ($P < .01$) than with SOY or SOY + FAT supplements.

Trial 2. Plasma glucose concentration (Table 3) was lower ($P < .1$) for control than for other S groups. Glucose concentration was higher ($P < .05$) at 9 wk than at 4 wk prepartum for all S groups (Table 3).

Neither albumin nor total protein was affected by supplement, but both were lower ($P < .01$) at 4 and 1 wk than at 9 wk prepartum (Table 3). Urea nitrogen concentration was lower ($P < .01$) in control than in other S groups. Concentration of urea nitrogen was higher ($P < .05$) for SOY + BM and SOY + CGM than for SOY and SOY + FAT. Creatinine was similar between treatments and weeks prior to parturition.

Supplement affected ($P < .01$; Table 3) blood serum cholesterol concentration. Cholesterol concentration was higher ($P < .01$) for SOY + FAT than for all other S groups.

At 9 wk prepartum, total bilirubin was lower than at 1 wk for all S groups. At 1 wk prepartum, bilirubin was lower for SOY + BM than for any other S groups (Figure 1).

Discussion

Supplementation with protein resulted in a positive response in BW change in both years. Cows receiving supplemental escape protein responded in both trials with reduced body condition loss and in Trial 2 with reduced

TABLE 3. LEAST SQUARES MEANS FOR PRECALVING CONCENTRATIONS OF GLUCOSE, ALBUMIN, TOTAL PROTEIN, UREA NITROGEN, CREATININE AND CHOLESTEROL AS INFLUENCED BY SUPPLEMENT AND WEEK PRIOR TO PARTURITION (TRIAL 2)

Metabolite	Supplement ^a					Week			
	Control	SOY	SOY+ BM	SOY+ CGM	SOY+ FAT	9	4	1	SE
Glucose mg/dl ^{bc}	54.0	56.1	57.0	56.0	57.7	58.6	55.4	54.1	1.6
Albumin, g/dl ^c	3.8	4.0	4.0	3.9	4.2	4.3	3.8	3.7	.3
Total protein, g/dl ^c	6.8	6.9	6.8	6.7	6.6	7.2	6.7	6.4	.2
Urea nitrogen, mg/dl ^d	7.1	10.5	12.7	12.5	10.3	9.9	10.5	11.4	1.0
Creatinine, mg/dl	1.8	1.9	1.8	1.8	1.9	1.8	1.9	1.9	.1
Cholesterol, mg/dl ^e	93.6	90.4	80.5	90.9	122.0	99.0	98.5	88.9	7.2

^aSOY = soybean meal, SOY+BM = soybean meal plus blood meal, SOY+CGM = soybean meal plus corn gluten meal, SOY+FAT = Soybean meal plus animal fat.

^bControl vs other supplements, $P < .10$.

^cEffect of week prior to parturition, $P < .05$.

^dControl vs other supplements, $P < .01$; SOY vs SOY+BM, SOY+CGM, $P < .05$; SOY+FAT vs SOY+BM, SOY+CGM, $P < .05$.

^eSOY+FAT vs SOY+BM, SOY+CGM, $P < .01$.

weight loss; in Trial 2, all cows were in greater negative energy balance compared to Trial 1 when cows gained BW. This response in Trial 2 is similar to that reported by Lindsay et al. (1982), who investigated a supplement containing both ruminally degraded and escape protein to a nonsupplemented control. They supplemented pregnant cows with a meat and bone meal, formaldehyde-treated cottonseed meal and fish meal supplement and obtained a positive weight change, whereas nonsupplemented cows lost BW. Differential response to similar supplement regimens in different years has been reported previously (Kartchner, 1981; Phillips and Vavra, 1981; Stanton et al., 1983). The appearance of a year effect in this study probably was not a function of ambient temperature. Beverlin (1988) noted that mean daily temperature did not differ between these two years. However, forage was covered by snow in Trial 2 for longer periods than in Trial 1. There also appeared to be less forage available in Trial 2, possibly due to a drought during the preceding growing season (National Oceanic and Atmospheric Administration, 1985, 1986). In both trials, forage always was available for grazing. However, initial grazing and selection may have reduced the quality of available forage in the second half of the trial in Trial 2 more than in Trial 1. Analysis of diet samples from both years showed similar ADF content of esophageal extrusa, although extrusa CP was higher in Trial 2 (Miner and Petersen, 1989).

Given the digestible OM intakes of this study as discussed by Miner and Petersen (1989), microbial protein synthesis can be estimated by using protein synthesis rates measured by Kropp et al. (1976) and Petersen et al. (1985). In turn, equations of NRC (1984) and Ørskov (1982) can be used to calculate a metabolic protein requirement of gestating beef cows. These estimations indicate that cows in the SOY group may have been deficient in metabolically available protein in




Figure 1. Trial 2 mean concentration of total bilirubin in gestating range cows. Periods represent sampling intervals in relation to weeks prior to parturition (Period 1, ○ = 9 wk; Period 2, ● = 5 wk and Period 3, ■ = 1 wk). Control = no supplement, SOY = soybean meal, SOY + BM = soybean meal plus blood meal, SOY + CGM = soybean meal plus corn gluten meal and SOY + FAT = soybean meal plus animal fat are the designations for supplement treatments. Supplement \times week prior to parturition interaction, $P < .01$, SE = .03.

both Trials 1 and 2. Because ruminal ammonia concentration was above 5 mg/100 ml of ruminal fluid (Miner and Petersen, 1989) in Trial 1, additional ammonia N in the rumen would not be expected to increase microbial protein yield (Satter and Slyter, 1974). Consumption of escape protein may increase the quantity of metabolically available protein in two ways. First, the escape protein should contribute directly to the pool of available protein. Microbial yield also may be enhanced because of enhanced availability of growth-limiting organic acids supplied by slowly ruminally degraded protein sources.

Alleviation of BW loss would be expected if the amino acids derived from escape protein substituted for or complemented amino acids of tissue origin in meeting amino acid requirements (NRC, 1984). An increase in amino acid pool size also may improve the efficiency of acetate utilization derived from either ruminal fermentation or adipose tissue catabolism (MacRae and Lobley, 1986). This suggested mechanism may be important in the subclinical ketosis experienced during weight loss and low plasma glucose concentrations noted in both trials. If amino acids of dietary origin accumulate, then those amino acids that serve as precursors of oxaloacetate may be utilized as intermediates for oxidation of acetate in the tricarboxylic acid cycle.

Another possible effect of escape protein involves the animal's requirement for essential amino acids. In Trial 1, serum amino acid concentrations were measured on samples composited across sampling intervals within period 2 (Miner, 1986). Branched-chain amino acid concentration was highest ($P = .07$) in SOY + BM and SOY + CGM (.63, .58, .52, .49 and .48 $\mu\text{mol/ml}$ for SOY + BM, SOY + CGM, SOY + FAT, SOY and control, respectively), which, according to Lynch and Jackson (1986), may reflect reduced muscle catabolism. Control (.08 $\mu\text{mol/ml}$), SOY (.07 $\mu\text{mol/ml}$) and SOY + FAT (.09 $\mu\text{mol/ml}$) had intermediate concentrations of lysine in comparison to SOY + BM (.11 $\mu\text{mol/ml}$), which was nearly double ($P < .05$) that of SOY + CGM (.06 $\mu\text{mol/ml}$). According to Ahmed (1982), who investigated methionine requirements in steers, SOY + CGM may have supplied a limiting essential amino acid that increased the use of lysine. For example, methionine may have been limiting because consumption of corn gluten meal did not affect

methionine concentration. Corn gluten meal protein is relatively rich in methionine but relatively low in lysine compared to blood meal (NRC, 1982). Lysine probably was not limiting because lysine concentration was elevated when lysine-rich blood meal was fed. However, these results and interpretation should be viewed with caution because we did not measure concentration of amino acids reaching the small intestine.

Weight change by cows fed SOY + FAT was intermediate compared to SOY + BM and SOY + CGM. The animal fat in SOY + FAT may have increased the amount of soybean meal that escaped ruminal degradation, although in situ measurements did not indicate this (Miner and Petersen, 1989). Lipids have been used to protect protein sources from fermentation (Van Soest, 1982). Even though the cattle consuming the SOY + FAT received twice the quantity of supplemental TDN, added fat did not alter weight change or improve nutritional status as measured by blood metabolite concentrations. From this, we may conclude that energy in the form of fat was not limiting to cow performance. The probability that forage intake (Miner and Petersen, 1989) was higher in SOY + BM than in SOY probably accounts for a portion of the observed differences in weight change in Trial 2.

During Trial 1, the lower glucose concentrations at 4 wk compared to 9 wk prepartum may have been a result of the environmental conditions. Snow covered much of the forage during 4 wk but was nearly absent during 9 wk and 1 wk. Snow cover reduces forage availability and may reduce digestibility, the effect being greater for unsupplemented heifers than for heifers fed a soybean meal supplement (Rittenhouse et al., 1970). In addition, mean ambient daily temperature was lower during 4 wk (-6.5°C) than for either 9 wk (-4.2°C) or 1 wk (1.1°C). Glucose turnover and oxidation are increased dramatically in sheep during acute cold exposure; skeletal muscles oxidize glucose for thermogenesis and the glucose supply is increased, probably due to increased glycogenolysis and gluconeogenesis (Sasaki and Weekes, 1986). Nevertheless, plasma glucose concentration has been shown to increase during cold exposure in most, but not all, cases (Sasaki and Weekes, 1986). Hence, our results may reflect the combined effect of snow cover and cold.

During Trial 2, glucose concentration in all 5 groups declined from 9 wk to 4 wk prepartum. This may reflect an increased fetal requirement for glucose and glucogenic substrate and is consistent with the linear decline in glucose with gestation observed by Prior and Scott (1977), Ferrell and Ford (1980) and Bull et al. (1984). Regulation of glucose concentration in the beef cow may be such that it falls below 50 mg/dl only in extreme situations.

Serum urea N concentration was higher in supplemented groups than in the control group in Trial 2 and higher in SOY + BM and SOY + CGM than SOY in both trials. This probably reflects the increased N supply of these diets and that most of the additional amino acids absorbed were deaminated by the liver and/or fetus. An increased amino acid supply should increase the supply of glucogenic intermediates.

Albumin concentrations, an indication of protein status, in Trial 1 were below, and in Trial 2 were above, the normal range reported by Benjamin (1978). The decline with advancing gestation in Trial 2 is consistent with results of Bull et al. (1984).

Total bilirubin concentration, an indication of liver function during physiological distress, was much higher in Trial 2 when most cows lost BW than in Trial 1 when most cows gained weight. In both trials, the concentration was within the normal range reported by Benjamin (1978). Concentration increased with advancing gestation in both trials, which is consistent with results of Bull et al. (1984), who found bilirubin to be increased by protein restriction in prepartum heifers. In Trial 2, SOY + BM had lower bilirubin than other treatments during 1 wk, indicating that cows in SOY + BM may have had a greater ability to cope with the demands of advancing gestation. Because bilirubin was higher during 4 wk than 1 wk prepartum of Trial 1, the hypothesis concerning environmental stress in 4 wk is supported.

Serum creatinine concentration, an indication of skeletal tissue catabolism, was higher in control than in other treatments in Trial 1; this is consistent with the response reported by Bull et al. (1984) to protein restriction. Creatinine concentration increased with advancing gestation, which also is consistent with results reported by Bull et al. (1984). This response in Trial 1 indicates that creatinine is

more sensitive to factors associated with advancing gestation than to the environmental stress of 4 wk.

The SOY + FAT treatment elevated serum cholesterol concentration. Elevated cholesterol can be indicative of dietary lipid content or tissue catabolism. Increased lipid intake has been shown previously to elevate cholesterol (Talavera et al., 1985). With advancing gestation, cholesterol has been shown to decline in dairy cows (Blum et al., 1983) but to increase in beef heifers (Bull et al., 1984). Our study showed a decline in 1 wk prepartum compared to 9 wk and 4 wk in both years.

Even during Trial 1, when most cows gained BW, certain blood metabolites (glucose, creatinine, bilirubin) reflected nutritional stress in unsupplemented and supplemented animals. Body weight change was determined precalving. In both trials, subtracting fetal and placental weight from final cow weight would reveal that most cows were losing BW during these trials.

When supplemented with ruminally degraded protein, the addition of ruminal escape protein may provide enhanced resistance to environmental and physiological (pregnancy) stress, as demonstrated by the consistent responses measured by improved body condition change and lower total bilirubin concentrations. This study also suggests that supplemental escape protein may reduce body condition and weight loss in years when unsupplemented cows would lose 50 kg in a 75-d period. Further studies are required to understand the role that escape protein may have for the range beef cow in negative energy balance.

Literature Cited

- Ahmed, B. M. 1982. Plasma amino acid studies in growing cattle. Ph.D. Dissertation. Michigan State Univ., East Lansing.
- Benjamin, M. M. 1978. Outline of Veterinary Clinical Pathology (3rd Ed.). Iowa State Univ. Press, Ames.
- Beverlin, S. K. 1988. Winter foraging behaviors of beef cows in relation to acute thermal stresses and supplementation. M.S. Thesis. Montana State Univ., Bozeman.
- Blum, J. W., P. Kunz and H. Leuenberger. 1983. Thyroid hormones, blood plasma metabolites and haematological parameters in relationship to milk yield in dairy cows. Anim. Prod. 36:93.
- Bull, R. C., D. O. Everson, D. P. Olson, L. F. Woodard and L. C. Anderson. 1984. Blood metabolite levels in pregnant beef heifers restricted prepartum in protein and/or energy. Proc. West. Sec. Am. Soc. Anim. Sci. 35:234.
- Ferrell, C. L. and S. P. Ford. 1980. Blood flow, steroid

- secretion and nutrient uptake of the gravid bovine uterus. *J. Anim. Sci.* 50:1113.
- Kartchner, R. J. 1980. Effects of protein and energy supplementation of cows grazing native winter range forage on intake and digestibility. *J. Anim. Sci.* 51:432.
- Kropp, J. R., R. R. Johnson, J. R. Males and F. N. Owens. 1977. Microbial protein synthesis with low quality roughage rations: Isonitrogenous substitution of urea for soybean meal. *J. Anim. Sci.* 45:837.
- Lindsay, J. A., G. W. J. Mason and M. A. Toleman. 1982. Supplementation of pregnant cows with protected proteins when fed tropical forage diets. *Proc. Aust. Soc. Anim. Prod.* 14:67.
- Lusby, K. S. and D. G. Wagner. 1987. Effects of supplements on feed intake. In: F. N. Owens (Ed.) *Feed Intake by Beef Cattle: Symposium*. Oklahoma State Univ. MP-121, Stillwater.
- Lynch, G. P. and C. Jackson, Jr. 1983. Metabolic responses of ewes to different protein intakes during late gestation. *Can. J. Anim. Sci.* 63:595.
- MacRae, J. C. and G. E. Lobley. 1986. Interactions between energy and protein. In: L. P. Milligan, W. L. Grovum and A. Dobson (Ed.) *Control of Digestion and Metabolism in Ruminants*, pp 367-385. Prentice-Hall, Englewood Cliffs, NJ.
- Mies, W. L., O. O. Thomas and C. W. Newman. 1967. Evaluation of biuret in cattle wintering and fattening rations. *J. Anim. Sci.* 26:925.
- Miner, J. L. 1986. Bypass supplementation of grazing pregnant beef cows. M.S. Thesis. Montana State Univ., Bozeman.
- Miner, J. L. and M. K. Petersen. 1989. The effects of ruminal escape protein or fat on ruminal characteristics of pregnant winter-grazing beef cows. *J. Anim. Sci.* 67:2782.
- National Oceanic and Atmospheric Administration. 1985. Climatological Data - Montana. National Oceanic and Atmospheric Administration, National Environmental Satellite Data and Information Service, Asheville, NC.
- National Oceanic and Atmospheric Administration. 1986. Climatological Data - Montana. National Oceanic and Atmospheric Administration, National Environmental Satellite Data and Information Service, Asheville, NC.
- NRC. 1982. United States-Canadian Tables of Feed Composition (3rd Rev. Ed.). National Academy Press, Washington, DC.
- NRC. 1984. Nutrient Requirements of Beef Cattle (6th Rev. Ed.). National Academy Press, Washington, DC.
- Ørskov, E. R. 1982. Protein Nutrition in Ruminants. Academic Press, New York.
- Petersen, M. K., and D. C. Clanton. 1981. Bypass protein (blood meal) in winter range supplements for grazing beef cow and weaned calf. *Proc. West. Sec. Am. Soc. Anim. Sci.* 32:18.
- Petersen, M. K., D. C. Clanton and R. Britton. 1985. Influence of protein degradability in range supplements on abomasal nitrogen flow, nitrogen balance and nutrient digestibility. *J. Anim. Sci.* 60:1324.
- Phillips, R. L. and M. Vavra. 1981. The effect of precalving energy levels on cow performance. *Proc. West. Sec. Am. Soc. Anim. Sci.* 32:117.
- Pinney, D. O., D. F. Stephens and L. S. Pope. 1972. Life time effects of winter supplementation feed level and age at first parturition on range beef cows. *J. Anim. Sci.* 34:1067.
- Prior, R. L. and R. A. Scott. 1977. Ontogeny of gluconeogenesis in the bovine fetus: Influence of maternal dietary energy. *Dev. Biol.* 58:384.
- Rittenhouse, L. R., D. C. Clanton and C. L. Streeter. 1970. Intake and digestibility of winter-range forage by cattle with and without supplements. *J. Anim. Sci.* 31:1215.
- SAS. 1984. SAS User's Guide: Statistics. SAS Inst., Inc., Cary, NC.
- Sasaki, Y. and T. E. C. Weekes. 1986. Metabolic responses to cold. In: L. P. Milligan, W. L. Grovum and A. Dobson (Ed.) *Control of Digestion and Metabolism in Ruminants*, pp 326-343. Prentice-Hall, Englewood Cliffs, NJ.
- Satter, L. D. and I. L. Slyter. 1974. Effect of ammonia concentration on rumen microbial protein production in vitro. *Br. J. Nutr.* 32:199.
- Stanton, T. L., F. N. Owens and K. S. Lusby. 1983. Formaldehyde-treated soybean meal for ruminants grazing winter range grass. *J. Anim. Sci.* 56:6.
- Talavera, F., C. S. Park and G. L. Williams. 1985. Relationships among dietary lipid intake, serum cholesterol and ovarian function in Holstein heifers. *J. Anim. Sci.* 60:1045.
- Thomas, O. O., J. L. Van Horn and R. L. Blackwell. 1968. Biuret and urea in supplements fed cattle grazed on native range. *Feeders Day Report PR-53*. Montana State Univ., Bozeman.